

References

1. Goletz, T.H., Henry, J.H., Continuous passive motion after total knee arthroplasty. *So. Med. Jo.* 79(9):1116, 1986.
2. Maloney, W.J., Schurman, D.J., Hangen, D., Goodman, S.B., Edworthy, S., Bloch, D.A. The influence of continuous passive motion on outcome in total knee arthroplasty. *Clin. Orthop.* 256:162, 1990.
3. Pope, R.O, Corcoran, S., McCaul, K., Howie, D.W. Continuous passive motion after primary total knee arthroplasty. Does it offer any benefits? *J. Bone Jt. Surg.* 79(6):914, 1997.
4. Worland, R.L., Arredondo, J., Angles, F., Lopez-Jimenez, F., Jessup, D.E. J. Home continuous passive motion machine versus professional physical therapy following total knee replacement. *J. Arthroplasty* 13(7):784, 1998.
5. Chen, B., Zimmerman, J.R., Soulen, L., De Lisa, J.A. Continuous passive motion after total knee arthroplasty: a prospective study. *Am. J. Phys. Med. Rehab.* 79(5):421, 2000.
6. Milne, S., Brosseau, L., Robinson, V. Noel, M.J., Davis, J. Drouin, H., Wells, G., Tugwell, P. Continuous passive motion following total knee arthroplasty. *Cochrane Database Syst. Rev.* 2:CD004260, 2003.